


Mesenchymal stem cell-derived extracellular vesicles attenuate multiple inflammasome pathways activated by chronic ethanol exposure in female mice

Blanca Martín-Urdiales and María Pascual


Department of Physiology, School of Medicine and Dentistry, University of Valencia, 46010, Valencia, Spain

BACKGROUND

Chronic alcohol consumption induces persistent neuroinflammation and inflammasome activation, contributing to alcohol-related brain damage. NLRP3 is one of the cytosolic sensors responsible for this activation through the formation of the NLRP3 inflammasome, which activates caspase-1 and promotes the release of inflammatory cytokines (e.g., IL-1 β , IL-18). Mesenchymal stem cell-derived extracellular vesicles (MSC-EVs) attenuate ethanol-induced NLRP3 activation, but their effect on other inflammasome pathways is unknown. We evaluated adipose-derived MSC-EVs on inflammasome-related gene expression in the prefrontal cortex, hippocampus, and striatum of female mice chronically exposed to ethanol.




METHODS



OBJETIVE: To evaluate whether adipose-derived MSC-EVs modulate multiple inflammasome pathways in the prefrontal cortex, hippocampus, and striatum of female mice chronically exposed to ethanol.

RESULTS

Chronic ethanol increased expression of NLRP3 pathway genes (*Nlrp3*, *Casp-1*, *Il1b*, *Il18*), upstream regulators (*Casp-11/4*, *Tlr4*), and alternative sensors (*Nlrp1*, *Aim2*, *Nlrc4*) across all three regions. MSC-EV treatment significantly attenuated these increases, restoring expression toward control levels. *Irak1* was unchanged.



CONCLUSIONS

Alcohol-induced neuroinflammation involves multiple innate immune pathways beyond NLRP3. MSC-EVs modulated inflammasome sensors, upstream regulators, and effector cytokines in regions key for cognition and reward, suggesting a broad immunomodulatory effect. MSC-EVs are a promising strategy against inflammasome-driven neuroinflammation in chronic alcohol consumption.

TAKE-HOME MESSAGE

MSC-EVs broadly **attenuate chronic ethanol-induced inflammasome activation** across brain regions key for cognition and reward, supporting their potential as a multifaceted therapy against alcohol-related neuroinflammation.